arXiv:1209.5847v1 [gr-qc] 26 Sep 2012

String Cosmology with Magnetic field in Anisotropic Space-time

C. P. Singh[1] **and Vijay Singh**[2]

Department of Applied Mathematics,
Delhi Technological University (Formerly Delhi College of Engineering)
Bawana Road, Delhi-110 042, India.
`cpsphd@rediffmail.com`[1]
`gtrcosmo@gmail.com`[2]

Abstract In this paper we study the effect of the magnetic field in string cosmology for a spatially homogenous and anisotropic Bianchi type -V space-time model. In order to study the effect of magnetic field, the standard form of the energy momentum tensor for cosmic strings is modified by including an additional term for magnetic field. The magnetic field is due to an electric current produced along the x-axis with infinite electrical conductivity. The field equations are solved for different string models such as geometric string (Nambu string), Takabayashi string (p-string) and Reddy string using string equation of state. We also find the solution for string models with uniform energy density. The physical and geometrical properties of each string model with and without magnetic field are discussed in detail.

Keywords: Cosmology; Anisotropic models; Massive string cosmology; Magnetic field.

PACS number(s): 98.80-k, 98.80-cq, 04.20-q.

1. Introduction

The search for cosmological models that do not suffer from the problems inflicting the standard big bang model of cosmology has intensified since the advent of string theory. String theory is particularly relevant to the initial singularity problem whose solution has long been thought to require a quantum theory of gravity, for which string theory seems to be the most promising candidate. It is generally believed that the very early universe underwent phase transitions, which gave rise to topological stable structures. Among the various topological defects that occurred during the phase transition and before the creation of particles in the early universe, strings have interesting cosmological consequences and have been studied in more details [1]. It is thought that cosmic strings cause density perturbations leading to the formation of galaxies [2]. These cosmic strings have stress-energy and couple with the gravitational field. The pioneering work in the formulation of the energy momentum tensor for the classical massive strings was done by Letelier [3], who considered the massive strings to be formed by geometric strings with particles attached along its extension. Stachel [4] developed a classical theory of the geometric strings. Letelier [5] first used this idea in obtaining some cosmological solutions of massive strings in Bianchi type-I and Kantowski - Sachs space-times. The gravitational effects of cosmic strings have been obtained by Vilenkin [6], Gott [7] and Garfinkle [8]. As the gravitational is the only long range force binding the contents of the universe and Einstein's theory of gravitation is the sole theory for understanding the nature and evolution of the large scale structure of the universe, so it may be interesting to study the gravitational effects which arise from strings within the framework of Einstein gravity. In literature, homogeneous and anisotropic cosmological models have been widely studied in string cosmology in different contexts (Ref. [9-21]).

On the other hand, the occurrence of magnetic field on galactic scale is well established fact today, and their importance for a variety of astrophysical phenomena is generally acknowledged. Several authors [22-28] have pointed out the importance of magnetic field in different context. The importance of the magnetic field for various astrophysical phenomenon has been studied in many papers. The string cosmological models with magnetic field have also been discussed by Banerjee et al. [9], Chakraborty [29], Tikekar and Patel [30, 31], Patel and Maharaj [32], Singh and Singh [33], Pradhan and Bali [34], and Bali [35]. Bali and Anjali [36] have investigated Bianchi Type-I magnetized string dust cosmological model. Bali et al. [37] have studied Bianchi-I massive string cosmological models with magnetic field. Wang [38], Pradhan et al.[39, 40], Pradhan [41, 42], Amirhashchi et al. [43], and Saha and Visinescu [44] have investigated the string cosmology with an incident magnetic field. Saha et al.[45] and Rikhvitsky et al. [46] have studied Bianchi-I string cosmological model in the presence of magnetic field and have examined the quantum effects in framework of loop quantum cosmology. Recently Singh [47] has generalized the work of Saha and Visinescu [44] to include the viscous term and has discussed the effect of viscous fluid with and without magnetic field in different string models for Bianchi I space -time.

Bianchi V universes are the natural generalization of Friedmann -Robertson- Walker (FRW) models with negative curvature. Bianchi V cosmological model where the matter moves orthogonally to the hypersurface of homogeneity, has been studied by Heckmann and Schucking [48]. Roy and Singh [49] have investigated Bianchi V model with stiff matter and a source free electromagnetic field. Banerjee and Sanyal [50] have studied Bianchi V model with viscosity and heat flow. Coley [51] has investigated Bianchi V imperfect fluid cosmological models in general relativity. Bali and

Meena [52] have discussed Bianchi V titled model for stiff matter. Chakraborty and Chakraborty [53], Bali and Singh [53], and Bali [54] have studied Bianchi V models with magnetic field in string cosmology. Singh et al.[54-58], Singh and Beesham [59] and Singh [60] have studied Bainchi V space time in general relativity with different contexts.

The purpose of this paper is to investigate a Bianchi type-V string cosmological model in the presence of magnetic field. The inclusion of the magnetic field is motivated by the observational cosmology and astrophysics indicating that many subsystems of the universe possess magnetic fields. We examine geometric string (Nambu string), Takabayashi string (p-string) and Reddy string models using the string equation of state. We also discuss the string models with uniform energy density. Exact analytic solutions are obtained in each case except for Takabayashi string model. The behavior of each model with and without magnetic field together with geometrical and physical aspects are discussed in detail.

The paper is organized as follows: In Sect.2 the general features of string theory and the corresponding field equations with the magnetic field are presented. A general soultion is given in Sect.3. In Sect. 4 we present the exact solutions for different string models. A solution with uniform energy density is presented in Sect.5. Finally, In Sect.6 we summarize our results.

2. Model and Basic Equations

We consider the homogenous and anisotropic Bianchi type -V spacetime whose metric is given by

$$ds^2 = -dt^2 + R_1^2 dx^2 + R_2^2 e^{-2x} dy^2 + R_3^2 e^{-2x} dz^2, \tag{1}$$

where R_1, R_2, and R_3 are the functions of cosmic time t only and are the scale factors in anisotropic background.

The energy momentum tensor for a cloud string with a magnetic field is taken into the form (Bali,[54])

$$T_i^j = \rho u_i u^j - \lambda x_i x^j + E_i^j, \tag{2}$$

where ρ is the rest energy density for a cloud of strings with massive particles attached to them and is related to the strings tension density λ by the relation (Letelier, [3])

$$\rho = \rho_p + \lambda. \tag{3}$$

Here ρ_p is the rest energy density of the particles attached to the strings. The energy conditions [9](Strong, weak and dominant) demand $\rho \geq 0$, $\rho_p \geq 0$, while λ is unrestricted; it may be positive, negative or zero as well. Also u_i is the four velocity for the cloud of particles and x^i is the four vector representing the string's direction, i.e., the direction of anisotropy and we have

$$u_i u^i = -x_i x^i = -1, \;\; u_i x^i = 0 \;\; and \;\; x^i = \left(R_1^{-1}, 0, 0, 0\right). \tag{4}$$

In Eq. (2), E_{ij} is the electromagnetic field given by (Linchnerowicz [61])

$$E_i^j = \bar{\mu}\left[|h|^2\left(u_i u^j + \frac{1}{2} g_i^j\right) - h_i h^j\right], \tag{5}$$

where $\bar{\mu}$ is the magnetic permeability, and h_i the magnetic flux vector defined by

$$h_i = \frac{\sqrt{-g}}{2\bar{\mu}}\, \epsilon_{ijkl} F^{kl} u^j, \tag{6}$$

In the above F_{kl} is the electro-magnetic field tensor and ϵ_{ijkl} is the totally anti-symmetric Levi-Civita tensor density with $\epsilon_{0123} = 1$. Let us assume the coordinates to be comoving, so that $u^1 = 0 = u^2 = u^3$ and $u^4 = 1$. The incidental magnetic field is taken along x-axis, so that $h_1 \neq 0$, $h_2 = 0 = h_3 = h_4$. This leads to $F_{12} = 0 = F_{13}$ by virtue of (6). Also, $F_{14} = 0 = F_{24} = F_{34}$ due to the assumption of the infinite conductivity of the field (Maartens [62]). Hence the only non vanishing components of F_{ij} is F_{23}. Therefore, the first set of Maxwell's equations

$$F_{ij;k} + F_{jk;i} + F_{ki;j} = 0. \tag{7}$$

leads to

$$F_{23} = \text{constant} = I(say). \tag{8}$$

In Eq. (7) the semicolon stands for covariant differentiation. From (6), we get

$$h_1 = \frac{R_1 I}{\bar{\mu} R_2 R_3}. \tag{9}$$

Since

$$|h|^2 = h_l h^l = h_1 h^1 = g^{11}(h_1)^2,$$

Therefore,

$$|h|^2 = \frac{I^2}{\bar{\mu}^2 R_2^2 R_3^2}\ . \tag{10}$$

Finally, one finds the following non-trivial components for E_i^j

$$E_0^0 = -\frac{I^2}{2\bar{\mu} R_2^2 R_3^2} = E_1^1 = -E_2^2 = -E_3^3, \tag{11}$$

and for the components of the string's energy momentum tensor, we have

$$T_0^0 = -\rho,\ T_1^1 = -\lambda,\ T_2^2 = T_3^3 = 0 = T_j^i,\ \ (i \neq j) \tag{12}$$

The Einstein's field equations (in gravitational units $8\pi G/c^4 = 1$) read as

$$R_i^i - \frac{1}{2} R g_i^j = -T_i^j. \tag{13}$$

The field equation (13) with (2) for line element (1) subsequently lead to the following system of equations

$$\frac{\dot{R}_1 \dot{R}_2}{R_1 R_2} + \frac{\dot{R}_2 \dot{R}_3}{R_2 R_3} + \frac{\dot{R}_3 \dot{R}_1}{R_3 R_1} - \frac{3}{R_1^2} = \rho + \frac{I^2}{2\bar{\mu} R_2^2 R_3^2}, \tag{14}$$

$$\frac{\ddot{R}_1}{R_1} + \frac{\ddot{R}_3}{R_3} + \frac{\dot{R}_1 \dot{R}_3}{R_1 R_3} - \frac{1}{R_1^2} = -\frac{I^2}{2\bar{\mu} R_2^2 R_3^2}, \tag{15}$$

$$\frac{\ddot{R}_1}{R_1}+\frac{\ddot{R}_2}{R_2}+\frac{\dot{R}_1\dot{R}_2}{R_1R_2}-\frac{1}{R_1^2}=-\frac{I^2}{2\bar{\mu}R_2^2R_3^2}, \tag{16}$$

$$\frac{\ddot{R}_2}{R_2}+\frac{\ddot{R}_3}{R_3}+\frac{\dot{R}_2\dot{R}_3}{R_2R_3}-\frac{1}{R_1^2}=\lambda+\frac{I^2}{2\bar{\mu}R_2^2R_3^2}, \tag{17}$$

$$2\frac{\dot{R}_1}{R_1}-\frac{\dot{R}_2}{R_2}-\frac{\dot{R}_3}{R_3}=0, \tag{18}$$

where a dot denotes ordinary derivative with respect to cosmic time t.

We define the various physical parameters such as the mean Hubble parameter H, anisotropy parameter A and shear scalar σ as

$$H=\frac{1}{3}\left(H_1+H_2+H_3\right), \tag{19}$$

$$A=\frac{1}{3}\sum_{i=1}^{3}\left(\frac{H_i-H}{H}\right)^2, \tag{20}$$

$$\sigma^2=\frac{1}{2}\,\sigma_{ij}\sigma^{ij}=\frac{1}{3}\theta^2-\left(\frac{\dot{R}_1\dot{R}_2}{R_1R_2}+\frac{\dot{R}_2\dot{R}_3}{R_2R_3}+\frac{\dot{R}_3\dot{R}_1}{R_3R_1}\right). \tag{21}$$

where $H_1=\frac{\dot{R}_1}{R_1}$, $H_2=\frac{\dot{R}_2}{R_2}$ and $H_3=\frac{\dot{R}_3}{R_3}$ are the directional Hubble parameters in directions of x, y and z axes, respectively and the scalar expansion is defined by $\theta=3H$. Also, $\sigma_{ij}=u_{i;j}+\frac{1}{2}\left(u_{i;k}u^ku_j+u_{j;k}u^ku_i\right)+\frac{1}{3}\theta\left(g_{ij}+u_iuj\right)$.

3. Solution of the field equations

We follow the method used by Saha and Visineseu [44] to solve the field equations (14)-(18). From (15) and (16), we get

$$\frac{d}{dt}\left(\frac{\dot{R}_3}{R_3}-\frac{\dot{R}_2}{R_2}\right)+\left(\frac{\dot{R}_3}{R_3}-\frac{\dot{R}_2}{R_2}\right)\left(\frac{\dot{R}_1}{R_1}+\frac{\dot{R}_2}{R_2}+\frac{\dot{R}_2}{R_2}\right)=0. \tag{22}$$

Let us define a new time-dependent function $\tau(t)$ as

$$\tau=R_1R_2R_3=\sqrt{-g}\ , \tag{23}$$

which is indeed the volume scale factor of the Bianchi V space-time. Using (23) into (22) and simplifies, we get

$$R_3=d_1R_2\exp\left[k_1\int\frac{dt}{\tau}\right], \tag{24}$$

where d_1 and k_1 are constants of integration. Eq.(18) gives

$$R_1^2=k_2R_2R_3, \tag{25}$$

where k_2 is a constants of integration without loss of generality we can take unity. Now from Eqs. (14)-(17) and (25), we get

$$\frac{\ddot{\tau}}{\tau}=\frac{1}{2}\left[(3\rho+\lambda)+\frac{R_1^2 I^2}{\bar{\mu}\tau^2}\right]+\frac{6}{\tau^{2/3}}. \tag{26}$$

On the other, the energy conservation law $T^{j}_{i\,;j}=0$ gives

$$\dot{\rho}+\frac{\dot{\tau}}{\tau}\rho-\frac{\dot{R}_1}{R_1}\lambda=0. \tag{27}$$

From (23) - (25), we find the following forms of metric functions in terms of τ (Ref. [55])

$$R_1=\tau^{\frac{1}{3}} \tag{28}$$

$$R_2=\frac{1}{\sqrt{d_1}}\,\tau^{\frac{1}{3}}\exp\left[-\frac{k_1}{2}\int\frac{dt}{\tau}\right] \tag{29}$$

$$R_3=\sqrt{d_1}\,\tau^{\frac{1}{3}}\exp\left[\frac{k_1}{2}\int\frac{dt}{\tau}\right] \tag{30}$$

It is to be noted that if the integration constant k_1 is taken to be zero, then anisotropic character of the model is lost and it reduces to a FRW model. Using (28)- (30), Eqs. (26) and (27) take the form

$$\ddot{\tau}=\frac{1}{2}(3\rho+\lambda)\tau+\frac{k}{2}\,\tau^{-\frac{1}{3}}+6\tau^{\frac{1}{3}}, \tag{31}$$

$$\dot{\rho}+\left(\rho-\frac{1}{3}\lambda\right)\frac{\dot{\tau}}{\tau}=0, \tag{32}$$

where $k=\frac{I^2}{\bar{\mu}}$.

To find the value of τ in terms of t, let us consider the equation of state for a cloud of string models[6]

$$\rho=\alpha\lambda, \tag{33}$$

where the constant α is defined by

$$\begin{aligned}\alpha &= 1 \quad (geometric\ or\ Nambu\ string),\\ &= (1+\omega) \quad (p-\ string\ or\ Takabayasi\ string),\\ &= -1 \quad (Reddy\ string),\end{aligned} \tag{34}$$

where ω is a constant such that $\omega>0$. Using (33) into (32), we get

$$\frac{\dot{\rho}}{\rho}=\left(\frac{1-3\alpha}{3\alpha}\right)\frac{\dot{\tau}}{\tau}, \tag{35}$$

which is integrated to give the solution

$$\rho=\rho_0\tau^{\frac{1}{3\alpha}-1}, \tag{36}$$

where ρ_0 is a positive constant of integration.

Now, Eq. (31) takes the form

$$\ddot{\tau} = \frac{1}{2}\left(\frac{3\alpha+1}{\alpha}\right)\rho_0\tau^{\frac{1}{3\alpha}} + \frac{k}{2}\tau^{-\frac{1}{3}} + 6\tau^{\frac{1}{3}}, \tag{37}$$

whose first integral is given by

$$\dot{\tau}^2 = 3\rho_0\tau^{\frac{3\alpha+1}{3\alpha}} + \frac{3}{2}k\tau^{\frac{2}{3}} + 9\tau^{\frac{4}{3}} + c_0, \tag{38}$$

which may be rewritten as

$$\dot{\tau} = \sqrt{3\rho_0\tau^{\frac{3\alpha+1}{3\alpha}} + \frac{3}{2}k\tau^{\frac{2}{3}} + 9\tau^{\frac{4}{3}} + c_0}, \tag{39}$$

where c_0 is a constant of integration. It is to be mentioned that τ is non-negative due to the expansion of the universe. At the points where $\tau = 0$, there occurs space-time singularity. On the other hand, the radical in (39) should be positive. Taking into account the energy density and string tension density obeying equation of state (33), we conclude that ρ and λ, and hence the right hand side of Eq. (31) is a function of τ only, i.e.,

$$\ddot{\tau} = F(\tau). \tag{40}$$

From the mechanical point of view, Eq. (40) can be interpreted as equation of motion of a single particle with unit mass under the force $F(\tau)$. Then the following first integral exists

$$\dot{\tau} = \sqrt{2[\epsilon - u(\tau)]}, \tag{41}$$

where ϵ can be viewed as energy level and $u(\tau)$ is the potential of the force F. Comparing Eqs. (39) and (41), we find $\epsilon = c_0/2$ and

$$u(\tau) = -\frac{1}{2}\left[3\rho_0\tau^{\frac{3\alpha+1}{3\alpha}} + \frac{3}{2}k\tau^{\frac{2}{3}} + 9\tau^{\frac{4}{3}}\right]. \tag{42}$$

Finally, we write the solution to Eq. (39) in a general quadrature form as

$$\int \frac{d\tau}{\sqrt{3\rho_0\tau^{\frac{3\alpha+1}{3\alpha}} + \frac{3}{2}k\tau^{\frac{2}{3}} + 9\tau^{\frac{4}{3}} + c_0}} = t + t_0, \tag{43}$$

where the integration constant t_0 can be taken to be zero.

It is too difficult to integrate (43) to find the exact value of τ in terms of t. Therefore, we look the solutions for different string models viz. geometrical, Takabayashi and Reddy strings using string equation of state (34) and the solution with uniform energy density in Sections 4 and 5.

4. Solution with equation of state

Let us solve the system of equations for the following string models to discuss the nature of solutions with and without magnetic field.

4.1 Geometric string model($\alpha = 1$)

For $\alpha = 1$ and $c_0 = 0$, Eq. (43) reduces to

$$\int \frac{d\tau}{\sqrt{3(\rho_0+3)\tau^{\frac{4}{3}} + \frac{3}{2}k\tau^{\frac{2}{3}}}} = t, \tag{44}$$

which on integration it gives

$$\tau = (c_1 t^2 - c_2)^{3/2}, \tag{45}$$

where $c_1 = (1 + \frac{\rho_0}{3})$ and $c_2 = \frac{k}{2(\rho_0+3)} = \frac{I^2}{2\bar{\mu}(\rho_0+3)}$.

At $t = 0$, τ becomes imaginary. For reality of the model, t must satisfy $t > \sqrt{\frac{3k}{2(\rho_0+3)^2}}$. From (28)-(30) and (45), we get the following form of the scale factors in terms of t.

$$R_1 = \sqrt{c_1 t^2 - c_2}, \tag{46}$$

$$R_2 = \frac{1}{\sqrt{d_1}}\sqrt{c_1 t^2 - c_2}\ \exp\left[\frac{k_1 t}{2c_2\sqrt{c_1 t^2 - c_2}}\right], \tag{47}$$

$$R_3 = \sqrt{d_1}\sqrt{c_1 t^2 - c_2}\ \exp\left[-\frac{k_1 t}{2c_2\sqrt{c_1 t^2 - c_2}}\right]. \tag{48}$$

We observe that R_1 and R_3 tend to zero but R_2 becomes indeterminate at $c_1 t^2 - c_2 = 0$. It is to be noted here that c_2 is the term which contains the magnetic field. Figure 1. shows that R_1 and R_3 always increase with time whereas R_2 decreases first then gradually increases with time in the presence of magnetic field. As $t \to \infty$, R_1 and R_2 tend to infinity whereas R_3 tends to zero. In fig. 1 we have assumed $\rho_0 = 3$, $I = 1$, $\bar{\mu} = 1.00001$, $d_1 = 10^4$, $c_1 = 2$ and $k_1 = 1$. In the absence of magnetic field, i.e., $c_2 = 0 = I$, we find that R_1 varies linearly where as R_2 and R_3 become indeterminate forms.

The directional Hubble parameters along x, y and z axes are respectively given by



Fig. 1 Evolution of the universe in the presence of magnetic field

$$H_1 = \frac{c_1 t}{c_1 t^2 - c_2}, \tag{49}$$

$$H_2 = \frac{c_1 t}{c_1 t^2 - c_2} - \frac{k_1}{2(c_1 t^2 - c_2)^{\frac{3}{2}}}, \tag{50}$$

$$H_3 = \frac{c_1 t}{c_1 t^2 - c_2} + \frac{k_1}{2(c_1 t^2 - c_2)^{\frac{3}{2}}}. \tag{51}$$

The average Hubble parameter in terms of cosmic time t is

$$H = \frac{c_1 t}{c_1 t^2 - c_2}. \tag{52}$$

The anisotropic parameter and shear scalar, respectively have the following expressions

$$A = \frac{k_1^2}{6c_1^2 t^2 (c_1 t^2 - c_2)}, \tag{53}$$

$$\sigma^2 = \frac{k_1^2}{4(c_1 t^2 - c_2)^3}. \tag{54}$$

The deceleration parameter is given by

$$q = \frac{c_2}{c_1 t^2}. \tag{55}$$

One can observe that as $t \to \infty$, $q \to 0$, which shows the marginal inflation (Coasting cosmology) in late time evolution of the universe. The anisotropy parameter and shear scalar also tend to zero as $t \to \infty$. The expression for the ratio $\frac{\sigma}{\theta}$ gives

$$\frac{\sigma}{\theta} = \frac{k_1}{6c_1 t\sqrt{(c_1 t^2 - c_2)}}. \tag{56}$$

We find that as $t \to \infty$, (56) tends to zero, which shows that the model tends to be isotropic for large t. The energy density and string tension density are respectively calculated as

$$\rho = \lambda = \rho_0 (c_1 t^2 - c_2)^{-1}. \tag{57}$$

We observe that ρ and λ are decreasing function of time. As t increases, both the parameters gradually decrease and approach to zero as $t \to \infty$. Thus, the string tension density disappears in late times. In this case the particle density, ρ_p is zero. The potential in terms of t is written as

$$u(t) = -\frac{3}{4}\left[k(c_1 t^2 - c_2) + 2(3 + \rho_0)(c_1 t^2 - c_2)^2\right]. \tag{58}$$

Figure 2. illustrates the potential graph with and without magnetic field for $\rho_0 = 3$, $c_1 = 2$ and $\bar{\mu} = 1.00001$. One may observe that the potential starts from zero and goes to positive for some finite time of interval and then becomes negative in the presence of magnetic field whereas it starts from zero and remains negative through out the evolution in the absence of magnetic field.



Fig. 2 Potential versus time in presence (bold line) and in absence (dashed line) magnetic field.

4.2. Reddy string model ($\alpha = -1$)

For $\alpha = -1$ and $c_0 = 0$, Eq. (43) reduces to

$$\int \frac{d\tau}{9\tau^{\frac{4}{3}} + 3\left(\rho_0 + \frac{k}{2}\right)\tau^{\frac{2}{3}}} = t, \tag{59}$$

The integral of (59) gives

$$\tau = (t^2 - c)^{3/2}, \tag{60}$$

where $c = \frac{k+2\rho_0}{6}$. At $t = 0$, τ becomes imaginary. For realty of the model, t must satisfy $t > \sqrt{\frac{k+2\rho_0}{6}}$. From (28)-(30) and (60), we get

$$R_1 = \sqrt{t^2 - c}, \tag{61}$$

$$R_2 = \frac{1}{\sqrt{d_1}}\sqrt{t^2 - c}\ \exp\left[\frac{k_1 t}{2c\sqrt{t^2 - c}}\right], \tag{62}$$

$$R_3 = \sqrt{d_1}\sqrt{t^2 - c}\ \exp\left[-\frac{k_1 t}{2c\sqrt{t^2 - c}}\right]. \tag{63}$$

The directional Hubble parameters are given by

$$H_1 = \frac{t}{t^2 - c}, \tag{64}$$

$$H_2 = \frac{t}{t^2 - c} - \frac{k_1}{2(t^2 - c)^{\frac{3}{2}}}, \tag{65}$$

$$H_3 = \frac{t}{t^2 - c} + \frac{k_1}{2(t^2 - c)^{\frac{3}{2}}}, \tag{66}$$

where as the average Hubble parameter has the form

$$H = \frac{t}{t^2 - c}. \tag{67}$$



Fig. 3 Evolution of the universe in presence (bold lines) and in absence (dashed lines)of magnetic field.

The anisotropic parameter, shear scalar and deceleration parameter have the following expressions, respectively.

$$A = \frac{k_1^2}{6t^2(t^2 - c)}, \tag{68}$$

$$\sigma^2 = \frac{k_1^2}{4(t^2 - c)^3}, \tag{69}$$

$$q = \frac{c}{t^2}. \tag{70}$$

The expression for the ratio $\frac{\sigma}{\theta}$ gives

$$\frac{\sigma}{\theta} = \frac{k_1}{6t\sqrt{(t^2 - c)}}. \tag{71}$$

The energy density, string tension density and particle density have the following expressions, respectively.

$$\rho = \rho_0(t^2 - c)^{-2}, \tag{72}$$

$$\lambda = -\rho_0(t^2 - c)^{-2}, \tag{73}$$

$$\rho_p = 2\rho_0(t^2 - c)^{-2}. \tag{74}$$

The potential in terms of t is given by

$$u(t) = -\frac{3}{4}\left[k(t^2 - c) + 6(t^2 - c)^2 + 2\rho_0(t^2 - c)^{-1}\right]. \tag{75}$$

We observe from figure 3 that the scale factors have the same behaviors in the presence of magnetic field as illustrated in fig.1. In the absence of magnetic field as shown by dashed lines in fig.3, the scale factors start expanding with different time of evolution. We have used $\rho_0 = 3$, $I = 1$, $\bar{\mu} = 1.00001$ and $d_1 = 100$ in fig.3. The other physical parameters such as deceleration parameter, anisotropy parameter, shear scalar also have the similar behaviors as discussed in section 4.1. For large t, all these parameters tend to zero. Therefore, the model approaches isotropic in later stage of the evolution of the universe. The energy density and particle density remain positive



Fig. 4 Energy density, string tension density and particle density versus time in presence (bold lines) and in absence (dashed lines) of magnetic field.



Fig. 5 Potential versus time in presence (bold line) and in absence (dashed line) magnetic field.

and decrease with time, and tend to zero for large t. The string tension density λ is negative and gradually increases with time, and finally it approaches to zero as $t \to \infty$. It is also observed that $\rho_p > |\lambda|$, i.e., the particle density remains larger than the string tension density during the cosmic expansion, especially in the early universe. The behaviors of these three parameters and potential with and without magnetic field are shown in figures 4 and 5, respectively with $\rho_0 = 3$, $I = 1$ or 0 and $\bar{\mu} = 1.00001$. The potential shows both positive and negative characters with and without magnetic field in this string model.

4.3 Takabayashi string model $(\alpha = 1 + \omega)$

For $\alpha = 1 + \omega$ and $c_0 = 0$, Eq. (43) reduces to

$$\int \frac{d\tau}{\sqrt{3\rho_0 \tau^{\frac{4+3\omega}{3(1+\omega)}} + \frac{3}{2}k\tau^{\frac{2}{3}} + 9\tau^{\frac{4}{3}}}} = t, \tag{76}$$

One can observe that it is too difficult to find the general solution of τ in terms of t. Therefore, we express the energy density, string tension density, particle density and potential in terms of τ as

$$\rho = \rho_0 \tau^{-\frac{2+3\omega}{3(1+\omega)}}, \tag{77}$$

$$\lambda = \frac{\rho_0}{(1+\omega)} \tau^{-\frac{2+3\omega}{3(1+\omega)}}, \qquad (\omega > 0) \tag{78}$$

$$\rho_p = \frac{\omega}{1+\omega} \rho_0 \tau^{-\frac{2+3\omega}{3(1+\omega)}}, \tag{79}$$

$$u(\tau) = -\frac{3}{2}\left[\rho_0 \tau^{\frac{3\omega+4}{3(1+\omega)}} + \frac{3}{2} k \tau^{\frac{2}{3}} + 9\tau^{\frac{4}{3}}\right]. \tag{80}$$

We observe that the physical parameters ρ, λ and ρ_p are decreasing functions of time and tend to zero as $t \to \infty$. Fig.6 shows the potential corresponding to Takabayashi string model with and without magnetic field for $\omega = 1$, $\rho_0 = 3$, and $\bar{\mu} = 1.00001$. It is always negative through out the evolution of the universe with and without magnetic field and decreases fast in the presence of magnetic field.



Fig. 6 Potential versus time in presence (bold line) and in absence (dashed line) magnetic field.

5. Solution with uniform energy density

Now we consider the rest energy density of the cloud of strings is uniform (i.e., $\rho = \rho_1$). From (32), we have $\tau = \tau_0$=constant. Eq.(31) gives

$$\lambda = -(3\rho_1 + k\tau_0^{-4/3} + 12\tau_0^{-2/3}). \tag{81}$$

Using the relation (4), we get

$$\rho_p = 4\rho_1 + k\tau_0^{-4/3} + 12\tau_0^{-2/3}. \tag{82}$$

The scale factors have the following expressions

$$R_1 = \tau_0^{1/3}, \tag{83}$$

$$R_2 = \frac{1}{\sqrt{d_1}}\, \tau_0^{1/3} \exp\left[-\frac{k_1}{2\tau_0} t\right], \tag{84}$$

$$R_3 = \sqrt{d_1}\, \tau_0^{1/3} \exp\left[\frac{k_1}{2\tau_0} t\right]. \tag{85}$$

We observe that the model is non-singular with the uniform density as R_1 is constant through out the evolution. The other two metric coefficients R_2 and R_3 are in exponential form. R_2 decreases to zero and R_3 exponentially grows with time. As λ is negative, therefore, the geometric string and Takabayashi string models are not possible due to negativity of the string tension density. The solution describes Reddy string model. During evolution of the universe the energy is getting transfer from string tension to particle density and their sum is always equal to ρ_1 and $\rho_p \geq 0$.

6. Conclusion

We have studied the anisotropic Bianchi type -V string cosmological model in the presence of magnetic field. The field equations have been solved exactly for the geometrical and Reddy string models. In both models we have found that the physical parameters have the same type of behaviors such as the scale factors along x and z axes increase with time but the scale factor along y axis decreases first and then increases with time, the energy density, particle density, anisotropy parameter and shear scalar all are decreasing functions of time and tend to zero as $t \to \infty$. In geometrical string model the string tension density decreases with time whereas it gradually increases with time and tends to zero for large t in Reddy string model as it is negative. It is also negative in the model with uniform energy density. All the string models tend to isotropic for large t. We have illustrated the scale factors, energy density, particle density, string tension density and potential with time for these models with and without magnetic field in figures 1-5.

In case of Takabayashi string model we observe that it is very difficult to find the solution due to complicated quadrature form of τ. Therefore, we have shown the potential versus τ graphically for this model the presence of magnetic field in fig.6. The reason to illustrate figures with and without magnetic field is to show the role of magnetic field in different string models. We have also discussed the string models with uniform energy density and have found a non-singular solution. In this model the geometric string and Takabayashi string models are not possible due to negativity of the string tension density.

References

1. Kibble, T.W.B.: J. Phys. A: Math. Gen. **9**, 1378 (1976).
2. Zel'dovich, Ya. B.: Mon. Not. R. Astron. Soc. **192**, 663 (1980).
3. Letelier, P.S.: Phys. Rev. D **20**, 1294 (1979).
4. Stachel, J.: Phys. Rev. D **21**, 2171 (1980).
5. Letelier, P.S.: Phys. Rev. D **28**, 2414 (1983).
6. Vilenkin, A.: Phys. Rep. **121**, 263 (1985).
7. Gott, J.R.: Astrophys. J. **288**, 422 (1985).
8. Garfinkle, D.: Phys. Rev. D **32**, 1323 (1985).
9. Banerjee, A., Sanyal, A.K., Chakraborty, S.: Pramana-J. Phys. **34**, 1 1990.
10. Chakraborty, S.: Ind. J. Pure Appl. Phys. **29**, 31 1991.
11. Ilhami , Y., Tarhan, I.: Astrophys. Space sci. **240**, 45 (1996).
12. Wang, X.X.: Chin. Phys. Lett. **20**, 615 (2003).
13. Pradhan, A., Yadav, A.K., Singh, R.P., Yadav, V.K.: Astrophys. Space Sci. **312**, 145

(2007).
14. Bali, R., Pradhan, A.: Chin. Phys. Lett. **24** 585 (2007).
15. Reddy, D.R.K., Naidu, R.L., Rao, V.U.M.: Int. J. Theor. Phys. **46**, 1443 (2007).
16. Rao, V.U.M., Vinutha, T., Shanthi, M.V.: Astrophys. Space Sci. **314**, 213 (2008).
17. Pradhan, A.: Commun. Theor. Phys. **51**, 367 (2009).
18. Pradhan, A., Amirhashchi, H., Yadav, M.K.: Fizika B **18**, 35 (2009).
19. Rao, V.U.M., Vinutha, T.: Astrophys. Space Sci. **325**, 59 (2010).
20. Pradhan, A., Chouhan, D.S.: Astrophys. Space Sci. **331**, 697 (2011).
21. Mahanto, K.L., Biswal, S.K., Sahoo, P.K., Adhikary, M.C.: Int. J. Theor. Phys. **51**, 1538, (2012).
22. Harrison, E.R.: Phys. Rev. Lett. **30**, 188 1973.
23. Melvin, M.A.: Ann. New York Acad. Sci. **262**, 253 1975.
24. Asseo, E., Sol, H.: Phys. Rep. **6**, 148 1987.
25. Parley, R., Taylor, G.: Astrophys. J. **101**, 1623 1991.
26. Wolfe, A.M., Lanzetta, K., Oren, A.L.: Astrophys. J. **388**, 17 (1992).
27. Kulstrud, R., Cen, R., Ostriker, J.P., Ryu, D.: Astrophys. J. **380** , 481 (1997).
28. Barrow, J.D.: Phys. Rev. D **55**, 7451, (1997).
29. Chakraborty, S.: Int. J. Pure Appl. Phys. **29**, 31, (1980).
30. Tikekar, R., Patel, L.K.: Gen. Relativ. Grav. **24**, 397, (1992).
31. Tikekar, R., Patel, L.K.: Pramana J. Phys. **42**, 482, (1994).
32. Patel, L.K., Maharaj, S.D.: Pramana J. Phys. **47**, 1, (1996).
33. Singh, G.P., Singh, T.: Gen. Relativ. Grav. **31**, 371, (1999).
34. Pradhan, A., Bali, R.: Euro. J. Theor. Phys. **5**, 91 (2008).
35. Bali, R.: Euro. J. Theor. Phys. **5**, 105, (2008).
36. Bali, R., Anjali: Astrophys. Space Sci. **302**, 901 (2006).
37. Bali, R., Pareek, U.K., Pradhan, A.: Chin. Phys. Lett. **24**, 2455, (2007).
38. Wang, X.X.: Chin. Phys. Lett. **23**, 1702, (2006).
39. Pradhan, A., Otarod, S., Singh, S.K.: Chin. J. Phys. **45**, 504, (2007).
40. Pradhan, A., Amirhashchi, H., Zainuddin, H.: Chin. J. Theor. Phys. **50**, 56, (2011).
41. Pradhan, A.: Fizika B **16**, 205, (2007).
42. Pradhan, A.: Commun. Theor. Phys. **51**, 367, (2009).
43. Amirhashchi, H., Zainuddin, H., Pradhan, A.: Int. J. Theor. Phys. **50**, 2531, (2011).
44. Saha, B., Visinescu, M.: Astrophys. Sapce. Sci. **315**, 99, (2008).
45. Saha, B., Rikhvitsky, V., Visinescu, M.: Cent. Eur. J. Phys. **8**, 113 (2010).
46. Rikhvitsky, V., Saha, B., Visinescu, M.: Astrophys. Sapce. Sci. **339**, 371, (2012).
47. Singh, C.P.: Astrophys. Space Sci. in press, (2012).
48. Heckmann, O., Schucking, E.: In Gravitation: An Introduction to Current Research ed. Witten, L. (John Wiley, New York), 1962.
49. Roy, S.R., Singh, J.P.: Aust. J. Phys. **38**, 763 1985.
50. Banerjee, A., Sanyal, A.K.: Gen. Relativ. Grav. **20**, 103 1988.
51. Coley, A.A.: Gen. Relativ. Grav. **22**, 3 1990.
52. Bali, R., Meena, B.L.: Proc. Nat. Acad. Sci. India **75**, 273 2005.
53. Chakraborty , N.C., Chakraborty S.: Int. J. Mod. Phys. D **10**, 723 2001.
54. Bali, R., Singh, D.K.: Astrophys. Space Sci. **300**, 387 2005.

55. Bali, R.: Eur. J. Thoer. Phys. **5**, 105 2008.
56. Singh, C.P., Zeyauddin, M., ShriRam: Astrophys. Space Sci. **315**, 181 2008.
57. Singh, C.P., Zeyauddin, M., ShriRam: Int. J. Mod. Phys.A **23**, 2719 2008.
58. Singh, C.P., Zeyauddin, M., ShriRam: Int. J. Theor. Phys. **47**, 3162 2008.
59. Singh, C.P., Beesham, A.: Int. J. Mod. Phys.A **25**, 3825 2010.
60. Singh, C.P.: Braz. J. Phys. **41**, 323 2011.
61. Linchnerowicz, A.: Relativistic Hydrodynamics and Magneto Hydrodynamics, Benjamin, New York, p.13 (1967).
62. Maartens, R.: Pramana J. phys. **55**, 575 (2000).